

COMPOSITIONS AND METHODS FOR PROTECTING CULTIVATED  
PLANTS FROM HERBICIDAL INJURY

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to U.S. Provisional Application Serial No. 60/092,993 which was filed on July 16, 1998.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

None.

BACKGROUND OF THE INVENTION

(1) Field of the Invention

The present invention relates to compositions comprising a herbicide, or combinations of herbicides, with or without a safener, and a repellent adjuvant, wherein the repellent adjuvant modifies the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. In particular, the herbicide composition comprises a repellent adjuvant that is an aqueous solution of an alkyltrialkoxysilane such as methyltrimethoxysilane and a water soluble silane coupling agent such as N-(2-aminoethyl)-3-aminopropyltrimethoxysilane or an aqueous solution of an organosiliconate such as sodium methyl siliconate.

(2) Description of Related Art

Many herbicides will cause injury to certain crop plants when applied in amounts that are effective in controlling weed growth. The damage to crop plants can be particularly severe when the crop plant is in an early stage of development, which is precisely the time when control of weed growth is most important. For this reason many herbicides are unsuitable for controlling

weeds when crop plants are at particular stages of growth. Therefore, the inability to control weed growth results in lower crop yield and reduced crop quality because the weeds compete with the crop plant for nutrients, light and water. In an attempt to broaden the usefulness of various herbicides, various herbicide compositions have been developed which contain compounds known as "safeners", also referred to as "antidotes", "protectants" or "antagonists."

The concept of using safeners to enhance the tolerance of crops to herbicides was developed in the late 1940's and has led to the development of particular herbicidal compositions that have the ability to control weed growth but without adversely affecting the growth or yield of a particular crop plant. However, the identification of an antidote which safens a particular class of herbicide or mixture of herbicides is not a theoretical determination but must be determined empirically. This determination is performed by observing the complex interaction of many biological and chemical factors, including the type of herbicide, the weed species to be controlled, the crop plant to be protected from weed competition and herbicidal injury, the developmental stage of the crop plant, and the safening compound itself. Moreover, the safener and herbicide must possess physico-chemical properties which allow an environmentally acceptable and stable product to be prepared. Therefore, because the discovery of herbicide/safener compositions is so empirically based, the development of effective herbicide/safener combinations is an expensive and unpredictable undertaking.

Isoxaflutole, an isoxazole herbicide, is a new soil-applied herbicide for use in corn. Researchers have reported that isoxaflutole provides excellent preemergence control of several broadleaf and annual grass weed species at rates ranging from 53 g/ha to 158

g/ha in conventional tillage and no-tillage corn (Bhowmik and Probst, Weed Sci. Soc. Am. Abstr. 36: 13 (1996); Curvey and Kapusta, North Central Weed Sci. Soc. 51: 57-58 (1996); Geier and Stahlman, North Central Weed Sci. Soc. 52: 81 (1997); Luscombe et al., Proc. North Central Weed Sci. Soc. 59: 57-58(1994); Mosier et al., Proc. North Central Weed Sci. Soc. 50: 74 (1995); Obermeier et al., Proc. North Central Weed Sci. Soc. 50: 25 (1995); Simkins et al., Proc. North central Weed Sci. Soc. 50: 25 (1995); Veilleux et al., North central Weed Sci. Soc. 50: 75 (1995); Vrabel et al., Proc. North Central Weed Sci. Soc. 24-25 (1995); Wrucke et al., Proc. North Central Weed Sci. Soc. 52: 17(1997); and Young et al., Weed Sci. Soc. Am. Abstr. 38: 8 (1998)). The mode of action of isoxaflutole is the competitive inhibition of the 4-hydroxyphenylpyruvate dioxygenase enzyme (EC 1.13.11.27) (Pallett et al., Pestic. Biochem. Physiol. 62: 113-124 (1998); Viviani et al., Pestic. Biochem. Physiol. 62: 125-134 (1998)). Inhibition of this enzyme disrupts carotenoid biosynthesis causing a bleaching symptomology in susceptible species similar to herbicides that disrupt carotenoid biosynthesis by targeting the phytoene desaturase enzyme (Luscombe and Pallett, Pestic. Outlook, 29-32 (1996); and Pallett et al., Pestic. Sci. 50: 83-84 (1997)). While isoxaflutole has been shown to be an effective herbicide for controlling weeds, isoxaflutole also has been shown to cause injury to corn and other crop plants, especially when applied postemergent. For an example, see U.S. Patent No. 5,627,131 to Shribbs et al.

Therefore, there is a need for a method for applying a herbicide post-emergence to a wide variety of cultivated plants that does not depend on the identity of the herbicide or the use of a safener. Such a composition would not be injurious to the cultivated plant but would maintain its ability to control weeds. The present invention provides compositions and methods

for their use that satisfy this need.

SUMMARY OF THE INVENTION

5 The present invention provides compositions  
comprising a herbicide, or a combination of herbicides,  
with or without a safener, and a repellant adjuvant  
which modifies the surface properties of the composition  
so that retention of the composition on foliage of the  
cultivated plant is reduced. An important element of  
10 the composition is the repellant adjuvant. When the  
repellant adjuvant is mixed with the herbicide or  
combination of herbicides, with or without a safener, to  
form the composition of the present invention and  
applied to the plants with a sprayer, the emitted  
15 composition spray forms spherical particles which then  
bounce off the plant foliage. In this manner the  
repellant adjuvant prevents retention of the composition  
by the plant. Thus, herbicidal injury to the cultivated  
plant is reduced because the herbicide composition is  
20 directed to the soil. Therefore, the present invention  
is particularly useful when it comprises herbicides that  
exert their primary effects at the soil.


The present invention provides a composition  
for protecting cultivated plants comprising (a) at  
25 least one herbicide and (b) a repellent adjuvant for  
modifying the surface properties of the composition so  
that retention of the composition on foliage of the  
cultivated plant is reduced. The composition of the  
present invention further comprises compositions wherein  
30 the herbicide is selected from the group consisting of  
acetanilides, acetamides, acetolactate synthase  
inhibitors, isoxazoles, diketonitriles, triketonitriles  
dinitroanilines, triazines, substituted ureas,  
ethofumerates, isoxafen, oxodiazon, dithiopyr and  
35 combinations thereof. Further still, the composition of  
the present invention comprises a composition wherein  
the composition further comprises a safener. In

particular embodiments of the present invention, the  
safener is selected from the group consisting of MON  
4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-  
dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine,  
5 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-  
dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-  
diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-  
furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-  
5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-  
10 spirocyclohexy-N-dichloroacetyl oxazolidine, 4-  
(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine,  
3-[3-(dichloroacetyl)-2,2-dimethyl-5-  
oxalidiny]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapиро-  
(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-  
15 methyl-2-isoquinolyl)ethanone, cis/trans-1,4-  
bis(dichloroacetyl)-2,5-dimethylpiperazine, N-  
(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-  
bis(dichloroacetyl)-1,5-diazacyclononane, 1-  
(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations  
20 thereof.

The present invention further provides a  
method for reducing injury to cultivated plants, by the  
herbicide, by applying as a spray at least one herbicide  
with or without a safener in a composition with the  
25 repellant adjuvant which modifies the surface properties  
of the composition so that retention of the composition  
on foliage of the cultivated plant is reduced. The  
present invention provides a method for protecting  
cultivated plants, the composition comprising (a) a t  
30 least one herbicide and (b) a repellent adjuvant for  
modifying the surface properties of the composition so  
that retention of the composition on foliage of the  
cultivated plant is reduced. The present invention  
further provides a method for using a composition  
35 wherein the herbicide is selected from the group  
consisting of acetanilides, acetamides, acetolactate  
synthase inhibitors, isoxazoles, diketonitriles,

triketonitriles dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Further still, the present invention provides a method for using the composition wherein the composition further comprises a safener. In particular embodiments of the present invention, the safener is selected from the group consisting of 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

To make the composition of the present invention or to practice the method of the present invention, it is preferable that the repellent adjuvant be selected from the group consisting of an aqueous solution of an alkali metal organosiliconate and an aqueous solution of a water soluble siloxane solution. The organosiliconate having the formula:



wherein X denotes sodium or potassium, and R is methyl, ethyl, or propyl, and the ration of Si:X is about 1:1; and an aqueous solution of a water soluble coupling

agent and an alkyltrialkoxysilane selected from the group consisting of alkyltrialkoxysilanes with C1 to C6 alkyl groups on silicon and a blend of alkyltrialkoxysilanes each with a C1 to C6 alkyl groups on silicon. In a preferred embodiment, the repellent adjuvant is selected from the group consisting of an aqueous solution of sodium methyl silicate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane.

#### OBJECTS

Therefore it is an object of the present invention to provide herbicide compositions that can be applied postemergence which do not injure cultivated plants while maintaining the herbicide's ability to effectively control weeds.

These and other objects will become increasingly apparent through the following detailed description of the invention and examples.

#### DESCRIPTION OF THE DRAWINGS

Figure 1 is a photograph that shows the ability of the composition of the present invention comprising an aqueous solution of sodium methyl silicate to prevent corn injury. The corn plants were treated from left to right: untreated control, corn treated with isoxaflutole in combination with DUAL II (metolachlor and benoxacor), and corn treated with a composition of the present invention (a combination of isoxaflutole, DUAL II, and sodium methyl silicate). The sodium methyl silicate was applied at a rate 0.25%. The photograph shows the corn 8 days after treatment.

Figure 2 is a photograph that shows the ability of the composition of the present invention comprising an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane


to prevent corn injury. The corn plants were treated from left to right: untreated control, corn treated with a combination of isoxaflutole and DUAL II, and corn treated with a composition of the present invention (a combination of isoxaflutole, DUAL II, and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane). The aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane was applied at a rate 0.25%. The photograph shows the corn 8 days after treatment.

#### DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention provides a composition for protecting cultivated plants comprising a herbicide, or combination of herbicides, with or without a safener, and a repellent adjuvant for modifying the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. The herbicide comprising the present invention can be selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxadiazon, dithiopyr and combinations thereof. In particular, it is desirable that the herbicide or combination of herbicides exert their primary effects at the soil. The repellent adjuvant comprising the composition is a silicone-based aqueous solution that forms an emulsion when in combination with the herbicide, which has modifies properties. The emulsion forms spherical particles when sprayed from a sprayer. The spherical particles bounce off the foliage of the plant to the ground where the herbicide then exerts its effect. One type of repellent adjuvant suitable for use in the present invention is exemplified by an aqueous solution of the



organosiliconate having the formula:



5 wherein X denotes sodium or potassium, and R is methyl, ethyl, or propyl, and the ratio of Si:X is about 1:1. In a preferred embodiment of an adjuvant of this type, the organosiliconate is sodium methyl siliconate, potassium methyl siliconate, or a mixture thereof. In  
10 a most preferred embodiment, the aqueous solution of organosiliconate consists essentially of 32 weight percent of sodium methyl siliconate and 67 weight percent of water. A second type of repellent adjuvant suitable for use in the present invention is an aqueous  
15 solution of a water soluble silane coupling agent and an alkyltrialkoxysilane, the alkyltrialkoxysilane being selected from the group consisting of alkyltrialkoxysilanes with C1 to C6 alkyl groups on silicon and a blend of alkyltrialkoxysilanes each with  
20 a C1 to C6 alkyl groups on silicon, the alkyltrialkoxysilane and the silane coupling agent preferably being present in the aqueous solution in the mole ratio of between about 0.5:1.0 to about 3.0:1.0. In a preferred embodiment, the water soluble silane  
25 coupling agent is N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and the alkyltrialkoxysilane is methyltrimethoxysilane. In a most preferred embodiment, the methyltrimethoxysilane, the N-(2-aminoethyl)-3-aminopropyltrimethoxysilane, and the water are in an  
30 aqueous solution consisting of a weight percent ratio of 35.7:58.2:6.1. Therefore, the silicon-based aqueous solution comprising the present invention is selected from the group consisting of an aqueous solution of an alkali metal organosiliconate and an aqueous solution of  
35 a water soluble silane coupling agent and an alkyltrialkoxysilane.

The composition of the present invention can

further include a second herbicide, an enhancer, or an adjuvant, which increases the activity or absorption of the first herbicide. In particular, the second herbicide is an acetanilide herbicide wherein the acetanilide herbicide increases the activity or absorption of the first herbicide in the composition. An example of an acetanilide herbicide is metolachlor or a mixture of the herbicide metolachlor and the safener benoxacor. In a preferred embodiment of the invention, the composition comprises an isoxazole herbicide such as isoxaflutole, an activator herbicide such as metolachlor, and the repellent adjuvant. In a more preferred embodiment of the invention, the composition comprises an isoxazole herbicide such as isoxaflutole, an activator herbicide such as metolachlor which is in combination with a safener such as benoxacor, and the repellent adjuvant.

The activity of certain herbicides can be increased by compounds that enhance absorption of the herbicide. Therefore, the present invention further includes compositions that comprise a herbicide, the repellent adjuvant, and an enhancer adjuvant which is an oil based adjuvant. In particular, the oil-based adjuvant is selected from the group consisting of a crop oil concentrate, a free fatty acid, and an esterified and saponified oil. Examples of herbicides that are known to work well in the presence of an oil-based adjuvant are the herbicides selected from the group consisting of cyclohexanidiones, aryloxyphenoxy, imidazolinone, and sulfonylurea herbicides.

The present invention also relates to a method for protecting cultivated plants including applying a herbicide formulation that has herbicidal activity from soil, the improvement comprising using as the herbicidal formulation a homogenous aqueous dispersion of the composition comprising a herbicide, and a repellent adjuvant for modifying the surface properties the

composition so that retention of the composition on foliage of the cultivated plant is reduced. The herbicide in the method of the present invention is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. The repellent adjuvant is selected from the group consisting of an aqueous solution of an alkali metal organosiliconate and an aqueous solution of a water soluble silane coupling agent and an alkyltrialkoxysilane. In a preferred embodiment, the repellent adjuvant is selected from the group consisting of an aqueous solution of methyl siliconate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapiro(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-

(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

5 The composition can further comprise an acetanilide herbicide as a second herbicide, which increases the activity or absorption of the first herbicide. In particular, the acetanilide herbicide in the composition can be metolachlor or a mixture of the herbicide metolachlor and the safener benoxacor. In a preferred embodiment of the invention the first  
10 herbicide comprising the composition is an isoxazole herbicide such as isoxaflutole.

15 The present invention further provides an improved method for protecting cultivated plants including the step of applying a herbicide compound, postemergence to the cultivated plants, the improvement which comprises adding a repellant adjuvant to the herbicide compound so that when the compound is applied to the cultivated plants, the compound bounces off foliage of cultivated plants to the soil. In the  
20 improved method, the repellent adjuvant is selected from any one of the repellent adjuvants disclosed herein. In the improved method, the herbicidal formulation can further comprise an enhancement material which enhances the activity of the herbicide. Enhancement materials  
25 can among other things include a second herbicide that activates or potentiates the activity of the first herbicide, or an oil-based adjuvant, or combinations of both. In a preferred embodiment, the herbicide is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors,  
30 isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the preferred method  
35 can include a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-

di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-  
2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-  
dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-  
dichloroacetyl oxazolidine, N,N-diallyl-2,2-  
5 dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-  
dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-  
N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-  
dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-  
10 dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-  
(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-  
(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-  
dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-  
isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-  
2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-  
15 tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-  
diazacyclononane, 1-(dichloroacetyl)-1-  
azaspiro[4,4]nonane, and combinations thereof.

The present invention further provides a  
method for protecting cultivated plants without injuring  
cultivated plants, the steps comprising: (a) providing  
20 a herbicidal formulation comprising one or more  
herbicides admixed with a repellent adjuvant wherein the  
repellant adjuvant modifies the surface properties of  
the formulation thereby reducing retention of the  
formulation on foliage of the cultivated plants; (b) and  
25 applying the formulation to the cultivated plants  
wherein the formulation bounces off the foliage onto the  
soil wherein the formulation protects the cultivated  
plants without injuring the cultivated plants. In the  
30 method, the repellant adjuvant is any one of the  
repellant adjuvants disclosed herein. In the present  
invention, the herbicidal formulation can further  
comprise an enhancement material which enhances the  
activity of the herbicide. In particular embodiments of  
35 the present invention, the herbicide is selected from  
the group consisting of acetanilides, acetamides,  
acetolactate synthase inhibitors, isoxazoles,

diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally comprises a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

The present invention further provides a method for inhibiting a weed without injuring turfgrass, the steps comprising (a) providing a liquid dispersion of a herbicidal formulation comprising one or more herbicides admixed with a repellent adjuvant wherein the repellent adjuvant modifies the surface properties of the formulation thereby reducing retention of the formulation on foliage of the turfgrass; and (b) applying the formulation to the crop plant wherein the formulation bounces off the foliage onto the soil wherein the formulation inhibits growth of the weed. In the method for protecting a turfgrass, the repellent adjuvant is any one of the repellent adjuvants disclosed

herein. In the method, the herbicidal formulation can further comprise an enhancement material which enhances the activity of the herbicide. In particular embodiments of the method, the herbicide is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

The amount of herbicide comprising the composition of the present invention and used in the method of the invention varies according to a number of parameters including the cultivated plant to be protected, the weed species to be controlled, and the edaphic and climatic conditions prevailing. In general, a rate of application from about 10 to 210 grams per

hectare (g/ha) of herbicide is suitable, preferably 50 to about 158 g/ha. The rate of the repellent adjuvant in the composition can be from 0.25% to 1.0%, preferably at a rate of 0.5%.

5           According to general cultivation practices, herbicides are mixed in a tank and applied to the plants using a sprayer. The practitioner will mix various combinations of herbicides in the tank, and in some cases, will include a safener to ameliorate the  
10 herbicide's activity towards the plant to be protected. In practicing the present invention, the practitioner in addition to the mixture of herbicides, with or without a safener, in the tank will include the repellent adjuvant to make the composition of the present  
15 invention.

Cultivated plants within the meaning of the present invention includes any plant cultivated for food or ornamentation with the exception of weeds. The cultivated plants to be protected by the method of the  
20 present invention include crop plants of which corn, sugarcane, beans, rice, wheat, oats, sorghum, and a wide variety of vegetables such as tomatoes, and fruits such as strawberries are examples. In a preferred embodiment, the method of the invention is performed  
25 where the crop to be protected is corn (*Zea mays*), sorghum (*Sorghum halepense*), sorghum (*Sorghum bicolor*), soybean (*Glycine max*) or dry bean (*Phaseolus vulgaris* L.). Examples of other cultivated plants that can be protected from a herbicide or combination of herbicides  
30 according to the present invention are turfgrasses; flowering garden plants such as roses, tulips, carnations, orchids and the like; various herb plants such as parsley, sage, rosemary, and thyme; ornamental plants such as shrubs, holly, juniper, and spice plants.

35           Thus, the objective of the present invention is to protect cultivated plants from injury from preemergence herbicides applied postemergence to the



crop. Generally, these herbicides have activity from the soil, however it is not necessary that they have activity from the soil. The present invention results in decreased herbicide injury to the cultivated plant because it decreases foliar retention and adsorption of the herbicide by the cultivated plant. Prior to the present invention, the use of certain herbicide combinations as a postemergence application were precluded because the combination either caused injury to the cultivated plant, or caused injury to the cultivated plants because the cultivated plants were in a particularly sensitive stage at the time of the herbicide application. Therefore, many herbicide combinations cannot be used for a wide variety of cultivated plants. While the examples disclosed herein describe use of the present invention for postemergence herbicides, the present invention is not to be construed as being limited to postemergence herbicides. For example, it can be desirable to apply a preemergence pesticide to an existing turfgrass stand to kill germinating and emerging crabgrass seedlings. The present invention comprising a preemergence herbicide and a repellant adjuvant would direct the herbicide to the soil wherein the herbicide would be active.

In addition to soil activity, previous research has shown that isoxaflutole has foliar activity on a number of weed species (Sprague et al., Weed Sci. Soc. Amer. Abstr. 37: 5 (1997); Vrable et al., *ibid.*; and Young and Hart, Weed Sci. 46: 397-402 (1998)), possibly allowing for the use of postemergence applications of isoxaflutole for weed control. In fact, Sprague et al. (*ibid.*) reported that postemergence applications of isoxaflutole at 105 g/ha controlled common lambsquarters, common ragweed, redroot pigweed, and velvetleaf greater than 90% and when it was tank-mixed with metolachlor/benoxacor, the mixture controlled foxtail by greater than 80%. But this tank-mixture

greatly reduced corn tolerance when it was applied after the corn had emerged. The basis for this corn injury was the result of increased isoxaflutole retention and subsequent absorption. This result indicated that metolachlor/benoxacor may act similar to various spray adjuvants that increase herbicide retention and thereby facilitating its subsequent absorption. Spray adjuvants are normally added to foliar-applied herbicide spray solutions to maximize the effectiveness of the herbicide. These adjuvants usually exert this enhancing effect by increasing herbicide spray retention on the leaf surface and by increasing herbicide penetration into the plant cuticle. A major barrier in the retention of a herbicide is the surface tension of the spray droplets. Adjuvants such as non-ionic surfactants (NIS) and 28% urea ammonium nitrate (UAN) have been found to decrease the surface tension of spray droplets, which results in an increase in surface coverage of the spray solution (De Ruiter et al., Weed Sci. 38: 567-572 (1990); Stevens et al., Pestic. Sci. 38: 237-245 (1993)). However, there are apparently no adjuvants which can be used to modify the surface properties of a herbicide solution, which in turn decreases herbicide retention and, therefore, the herbicide absorption. The present invention provides herbicide mixtures containing adjuvants which function as repellant adjuvants because they modify the surface properties of the mixtures. It is theorized that the repellant adjuvants modify the surface property of the mixture by causing an increase in the surface tension of the mixture, which results in spray droplets of increased surface tension. Because of the increased surface tension, the herbicide spray droplets are not retained by the plant foliage. Thus, the present invention permits the postemergence application of the herbicide mixtures disclosed herein.

The herbicide isoxaflutole by itself is not

injurious to corn when applied preemergence to the corn. However, when isoxaflutole is applied to corn postemergence it causes injury to the corn. The injury is particularly severe when isoxaflutole is combined with metolachlor and benoxacor, and the combination is applied to corn plants in the spike, 2-leaf, or 4-leaf stage. The reason is that the metolachlor or any other acetanilide herbicide applied with isoxaflutole increases spray retention on the corn leaves, which ultimately increases the absorption of isoxaflutole or any other isoxazole by the corn plant. For example, DUAL II is an oily composition which when mixed with the herbicide forms a composition that facilitates absorption of the herbicide by the leaf. It is the increased absorption of the isoxazole herbicide that causes the injury to corn. Therefore, when acetanilide herbicides are used in combination with other herbicides, the acetanilide herbicide may increase the retention and absorption of the other herbicide by the plant. Herbicide absorption can also be enhanced when in combination with oil-based adjuvants such as crop oil concentrate, free fatty acids, and esterified and saponified oils. Examples of such herbicides whose absorption is enhanced by oil-based adjuvants are cyclohexanidiones, aryloxyphenoxy, imidazolinone, and sulfonylurea herbicides.

The present invention is an improvement because it involves including in the herbicide spray solution, a material that modifies the surface properties of the spray solution, which results in decreased adherence of the spray droplets to the plant leaves. This is particularly useful when the plant is at a vulnerable stage of growth. For example, a crop such as dry beans can be at a sensitive growth stage (unifoliate) at the time it is most desirable to apply an acetanilide herbicide such as dimethamid. The present invention decreases foliar absorption of the herbicide

by the bean leaves, yet allows the herbicide to exert its action from the herbicide that has reached the soil.

Therefore, the present invention is a herbicide and water repellant composition wherein the surface properties of the composition is modified by the water repellant compound therein. Because these materials modify the surface properties of the herbicide spray solution, the spray droplets form spheres which bounce off the plant foliage. The water repellant compounds suitable for modifying the surface properties to make the herbicide composition of the present invention include an aqueous solution of an organosiliconate and aqueous siloxane solutions. An example of an aqueous solution of an organosiliconate that modifies the surface properties of the herbicide spray composition is an aqueous solution of sodium methyl siliconate. Sodium methyl siliconate has been used as a water repellant treatment for surfaces (see U.S. Patent No. 5,780,412 to Scarbrough and references therein). An example of an aqueous siloxane solution that is suitable is an aqueous solution of a water soluble silane coupling agent and an alkyltrialkoxysilane that modifies the surface properties of the herbicide spray composition such as an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane. When these compounds are mixed with an aqueous solution, they form an emulsion. Siloxanes are described in U.S. Patent 3,294,725 to Findlay et al. which is hereby incorporated herein by reference and aqueous solutions consisting of water soluble silane coupling agents and alkyltrialkoxysilanes are described in U.S. Patents 5,051,129 and 5,073,195, both to Cuthbert et al. which are hereby incorporated herein by reference to teach compositions which are suitable as repellant adjuvants in the present invention and methods for making them. While it may appear that any silicone containing

compound may be suitable for making the compositions of the present invention, the inventors have discovered otherwise. For example, methyltrimethoxysilane and a phosphonate ester alkyl silicon are water soluble silicon compounds. However, neither of these compounds is effective at producing a composition according to the present invention.

In an embodiment of the present invention disclosed herein, the herbicidal composition comprises isoxaflutole (5-cyclopropyl isoxazol-4-yl-2-mesyl-trifluoromethylphenyl ketone); a mixture of the herbicide metolachlor (2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide) or acetochlor, and the safener benoxacor ((4-dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine), MON 13900 or dichlormid; and any one of the repellant adjuvants disclosed herein. In a preferred embodiment, the herbicidal composition comprises the isoxaflutole as BALANCE, the metolachlor and benoxacor mixture as DUAL II, and the repellant adjuvant selected from the group consisting of an aqueous solution of sodium methyl siliconate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyl-trimethoxysilane.

While the examples disclosed herein relate to the herbicide isoxaflutole, the present invention is not to be construed as being limited to the herbicide isoxaflutole. Examples of other herbicides which are encompassed by the present invention are nicosulfuron which is 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-N,N-dimethyl-3-pyridinecarboxamide - (ACCENT, Dupont, Wilmington, Delaware; isopropylamine salt, glyphosphate with adjuvants, (ACCORD, Monsanto Company, St. Louis, Missouri); primisulfuron which is methyl 2-[[[(4,6-bis(difluoromethoxy)-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoate (BEACON, Novartis, Greensboro, North Carolina); Chlorimuron which is ethyl 2-[[[(4-chloro-

6-methoxy-2-pyrimidinyl)amino]carbonyl] amino]  
sulfonyl]benzoate ethyl (CLASSIC, Du Pont, Wilmington,  
Delaware); Glufosinate-ammonium salt which is (2-amino-  
4-(hydroxymethylphosphinyl)butanoic acid (LIBERTY,  
5 AgrEvo, Wilmington, Delaware); Linuron which is N1-(3,4-  
dichlorophenyl)-N-methoxy-N-methylurea) (LOROX, Bayer,  
Kansas City, Kansas); Linuron and chlorimuron ethyl  
(LOROX PLUS, Dupont, Wilmington, Delaware);  
Thifensulfuron which is (methyl 3-[[[(4-methoxy-6-  
10 methyl-1,3,5-triazin-2-yl]carbonyl]amino] sulfonyl]-2-  
thiophenecarboxylate) (PINNACLE, Dupont, Wilmington,  
Delaware); Imazethapyr (PURSUIT, American Cyanamid,  
Princeton, New Jersey); glyphosate-isopropyl amine salt  
which is (N-(phosphonomethyl)glycine) (ROUNDUP, Monsanto  
15 Company, St. Louis, Missouri); ROUNDUP with surface  
components (phosphate esters and cationic tallow amines  
(ROUNDUP ULTRA, Monsanto Company, St. Louis, Missouri);  
imazaquin which is (2-[4,5-dihydro-4-methyl-4-(1-  
methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinoline  
20 carboxylic acid) (SCEPTER, American Cyanamid, Princeton,  
New Jersey); acetochlor which is HARNESS and SURPASS  
(available from Monsanto Company, and Zeneca Ag-Products  
Wilmington, Delaware, respectively); alachlor, which is  
2-chloro-2',6'-diethyl-N-(methoxymethyl)acetanilide and  
25 sold as LASSO (available from Monsanto Company); EPTC  
which is S-ethyl dipropylthiocarbonate and sold as  
ERADICANE (available from Zeneca Ag-Products);  
halosulfuron which is PERMIT and BATTALION (available  
from Monsanto company); EPIC which is isoxaflutole, and  
30 flufenacet (BAYFOE 5043) which is 4-fluoro-N-isopropyl-  
2-[[5-trifluoromethyl)-1,3,4-thiadiazol-  
2yl]oxy]acetamide (available from Bayer); and  
glyphosphate-trimethylsulfonium salt (N-  
(phosphonomethyl)glycine) (TOUCHDOWN, Zeneca Ag-  
35 Products).

In addition to the composition of the present  
invention containing a herbicide and the repellant

adjuvant, the present invention can further comprise activators, enhancers and safeners. Therefore, the present invention can further comprise a monosaccharide wherein the monosaccharide acts as an enhancer or potentiator for the herbicide in killing the weed without decreasing tolerance of the crop to the herbicide. Examples of such compositions are in U.S. Application Ser. No. 08/984,407 filed December 3, 1997 which is herein incorporated by reference. The present invention can further include oil-based adjuvants such as crop oil concentrate, free fatty acids, and esterified and saponified oils. The present invention can further include a safener which causes a reduction in injury to the crop plant without an unacceptable reduction in the herbicidal action. Examples of safeners encompassed by the present invention include benoxacor which is (4-dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine; dichlormid which is 2,2-dichloro-N,N-di-2-propenylacetamide (available from Zeneca, Inc.); MON 4660 which is available from Monsanto Company; R-29148 which is 2,2,5-trimethyl-N-dichloroacetyloxazolidine (available from Zeneca Ag-Products); R-25788 which is N,N-diallyl-2,2-dichloroacetamide (available from Zeneca Ag-Products); and MON 13900 which is 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine (available from Monsanto Company). Other safeners include 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-

(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof. Safeners are also disclosed in U.S. Patent 5,627,131 to Shribbs et al. which is hereby incorporated herein by reference. Examples of particular herbicide and safener combinations include DUAL II which consists of metolachlor and benoxacor (available from Novartis); SURPASS which consists of acetochlor and dichlormid (available from Zeneca Ag-Products); MON 8407 which consists of acetochlor and MON 4660 (available from Monsanto Company); ERADICANE which consists of EPTC and R-29148 (available from Zeneca Ag-Products); BATTALION which consists of a halosulfuron and MON 13900 (available from Monsanto Company); and MON 8411 which consists of acetochlor and MON 13900 (available from Monsanto Company). A particularly desirable combination of herbicide and safener is the herbicide acetochlor mixed with a safener selected from the group consisting of dichlormid, MON-13900 (flurilazole), R-29148, R-25788 (dichlormid), MON 4660 and combinations thereof.

The following examples are intended to promote a further understanding of the present invention.

#### EXAMPLE 1

Various silicone-based compounds were evaluated for ability to prevent herbicidal injury to the corn when used as repellent adjuvants in herbicide formulations. In previous experiments, it was shown that isoxaflutole in combination with metolachlor and benoxacor applied once corn had emerged caused significant corn injury. The basis of this severe injury is the retention and subsequent absorption of isoxaflutole in the foliar tissue of the corn plant, which is exacerbated by the metolachlor herbicide in the composition. Therefore, to address this problem, a number of silicon-based compounds, which were believed



to have repellent properties, were evaluated for the ability to prevent injury to corn plants. The compounds evaluated were an aqueous solution of 32 weight percent sodium methyl siliconate and 67 weight percent water ("SMS"), an aqueous solution of 35.7 weight percent N-(2-aminoethyl)-3-aminopropyltrimethoxysilane, 58.2 weight percent methyltrimethoxysilane, and 6.1 weight percent water ("AFS-MTMS"), methyltrimethoxysilane ("MTMS"), and phosphonate ester alkyl silicon ("PEAS"). These test adjuvants are commercially available from Dow Corning, Midland, Michigan. The test adjuvants were applied in combination with isoxaflutole or isoxaflutole in combination with metolachlor and the safener benoxacor, and the ability of the combination to reduce plant injury by inhibiting retention and subsequent absorption of the herbicide was evaluated.

Pioneer 3573 corn seeds (Pioneer 3573, a product of Pioneer Hi-Bred International, Inc., Des Moines, Iowa) were planted 2.54 cm deep, and velvetleaf and barnyardgrass seeds were planted 1.0 cm deep in 875 ml pots containing BACCTO professional greenhouse potting mix (a product of Michigan Peat Co., Houston, Texas). The seedlings were grown in a greenhouse maintained at 25°C  $\pm$  2°C. Natural sunlight was supplemented with light from sodium vapor lamps, which provided a total midday light intensity of 1,000  $\mu\text{mol m}^{-2} \text{s}^{-1}$  photosynthetic photon flux at plant height during a 16 hour photoperiod. Plants were watered daily and fertilized weekly with 50 ml of a water soluble fertilizer solution (400 ppm nitrogen, 400 ppm  $\text{P}_2\text{O}_5$ , and 400 ppm  $\text{K}_2\text{O}$ ).

Isoxaflutole was combined with three different rates (0.25%, 0.5%, and 1.0%) of the four test adjuvants, which were then applied to postemergence to 2-leaf (5 inch) corn plants. Generally, corn leaf stages are described as the number of visible leaves. Isoxaflutole at 105 g/ha and isoxaflutole tank-mixed

with 1.1 kg/ha of metolachlor/benoxacor were applied either alone or in combination with ACTIVATOR 90, a non-ionic surfactant (NIS) product of Loveland Industries Inc., Greeley, Colorado) at 0.25% (v/v). The various herbicide applications were made to corn at the 2-leaf (V1) growth stage.

The herbicide compositions were applied through an 8003 E flat fan nozzle (available from Spraying Systems Co., Wheaton, Illinois) delivering 234 L/ha at a pressure of 172 kPa (25 gallons/acre). Corn tolerance was evaluated 8 days after treatment (DAT) by visually evaluating the plants for bleaching and necrotic symptoms and also by measuring corn height (base of the plant to its crown). Visual corn injury ratings were based on a scale from 0 to 100, with 0 indicating no effect and 100 indicating plant death. Corn height was measured in cm and presented as a percent of the non-treated plants, with 0 indicating total reduction in plant height and 100 indicating height equal to the non-treated plants. All experiments were conducted twice as completely randomized designs with four replications. Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at  $\alpha=0.05$ . Statistical analysis indicated no experimental run interactions, so the data were combined and reported as the means of two experiments. Non transformed means are presented since arcsine and square root transformations did not alter the interpretation of the data.

The data for the experiments are presented in Tables 1 and 2, which show that isoxaflutole when applied to corn by itself did not significantly injure the corn. But when isoxaflutole was applied in combination with metolachlor/benoxacor, severe corn injury of 47% occurred and plant height was reduced by 48%. However, when the mixture of isoxaflutole and metolachlor/benoxacor was mixed with either SMS or AFS-

MTMS, this injury was reduced to less than 10%. In contrast, neither of the other silicon-based compounds, MTMS or PEAS, reduced injury to corn when added to the isoxaflutole or isoxaflutole and metolachlor/benoxacor mixture. It is interesting that the other silicon-based compositions were not effective in ameliorating the herbicide's affect on the corn, in particular MTMS. Therefore, this example shows that only the adjuvants, SMS and AFS-MTMS, decreased the retention and subsequent absorption of isoxaflutole when either adjuvant was in combination with metolachlor/benoxacor.

5

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Table 1.  
Isoxaflutole Injury to Corn in Greenhouse Trials

Treatment	Adjuvant Rates			
	0%	0.25%	0.5%	1.0%
-----(% injury to corn)-----				
Isoxaflutole	2			
Isoxaflutole + Activator 90		15		
Isoxaflutole + PEAS		2	0	0
Isoxaflutole + SMS		0	0	0
Isoxaflutole + AFS-MTMS		0	0	0
Isoxaflutole + MTMS		1	0	0
Isoxaflutole + Metolachlor <sup>a</sup>	47			
Isoxaflutole + Metolachlor + Activator 90		59		
Isoxaflutole + Metolachlor + PEAS		52	49	51
Isoxaflutole + Metolachlor + SMS		8	9	7
Isoxaflutole + Metolachlor + AFS-MTMS		8	3	2
Isoxaflutole + Metolachlor + MTMS		51	56	56
<sup>a</sup> The formulation metolachlor contained the herbicide safener benoxacor				

Table 2  
Corn Height as a Percent of Control, 8 DAT in Green  
House Trials

5	Treatment	Adjuvant Rates			
		0%	0.25%	0.5%	1.0%
		-----(% of control height)-----			
	Isoxaflutole	100			
	Isoxaflutole + Activator 90		96		
10	Isoxaflutole + PEAS		107	110	107
	Isoxaflutole + SMS		107	107	110
15	Isoxaflutole + AFS-MTMS		107	107	114
	Isoxaflutole + MTMS		103	110	107
	Isoxaflutole + Metolachlor <sup>a</sup>	62			
20	Isoxaflutole + Metolachlor + Activator 90		66		
	Isoxaflutole + Metolachlor + PEAS		62	69	69
25	Isoxaflutole + Metolachlor + SMS		100	100	100
30	Isoxaflutole + Metolachlor + AFS-MTMS		107	107	103
	Isoxaflutole + Metolachlor + MTMS		69	62	66
35	<sup>a</sup> The formulation metolachlor contained the herbicide safener benoxacor				

The results of Tables 1 and 2 for the various  
herbicide mixtures containing adjuvants added at a rate  
of 0.5% are presented in Table 3. Table 3 shows that  
injury to 2-leaf corn by isoxaflutole was reduced to 0  
only when SMS or AFS-MTMS was mixed with the herbicide.  
The table further shows that injury to corn caused by

isoxaflutole in combination with metolachlor/benoxacor was reduced from 47% to 8% and 3% by SMS and AFS-MTMS, respectively. The table also shows that isoxaflutole in combination with metolachlor/benoxacor reduced corn height by about 40% whereas adding either SMS or AFS-MTMS to the composition completely abrogated any affect the composition had on corn growth. The table clearly shows that the silicon-based compositions, PEAS and MTMS, were not useful as repellent adjuvants in herbicidal compositions. Therefore, this example demonstrates the present invention comprising either SMS or AFS-MTMS reduces herbicidal injury to corn when applied postemergence.

Table 3

Treatment <sup>a</sup>	Rate	Injury <sup>b</sup>		Height	
		Isoxaflutole <sup>c</sup>	Isoxaflutole + metolachlor <sup>d</sup>	Isoxaflutole	Isoxaflutole + metolachlor <sup>d</sup>
		-----%-----	-----%-----	-----% of control-----	-----% of control-----
Alone	0.25% v/v	2	47	100	64
+NIS	0.5% v/v	15	59	98	65
+PEAS	0.5% v/v	0	49	110	69
+SMS	0.5% v/v	0	8	106	99
+AFS-MTMS	0.5% v/v	0	3	107	105
+MTMS	0.5% v/v	0	56	108	62
LSD <sub>0.05</sub>		-----4-----	-----8-----		

<sup>a</sup>Treatments were applied to 2-leaf (12) corn

<sup>b</sup>Visual injury ratings and corn heights were evaluated 8 DAT

<sup>c</sup>Isoxaflutole was applied at 105 g/ha.

<sup>d</sup>The formulation of metolachlor contained the herbicide safener benoxacor and was applied at 1 l kg/ha

## EXAMPLE 2

The greenhouse experiments showed that adding SMS or AFS-MTMS to a herbicide mixture rendered the herbicide safe for use on cultivated plants without reducing the herbicide's effective against weeds. However, greenhouse experiments are performed under controlled conditions. Therefore, field experiments were performed to assess how the present invention would

perform under actual farm conditions.

Conventional tillage experiments were conducted in 1998 to evaluate the influence of SMS and AFS-MTMS, on corn tolerance and weed control from postemergence applications of isoxaflutole alone and in tank-mixture with metolachlor/benoxacor. Experiments were conducted at the Michigan State University Crop and Soil Science Research Farm at East Lansing, Michigan on Capac sandy clay loam soil (fine-loamy, mixed mesic Acric Ochraqualfs) containing 3.1% organic matter with a pH of 6.3 in 1998.

Tillage consisted of moldboard plowing in the fall prior to spring disking and field cultivation. Prior to spring cultivation, 320 kg/ha of 46-0-0 fertilizer was applied broadcast. At planting, 140 kg/ha of 6-24-24 fertilizer was applied as a banded treatment 5 cm below and 5 cm beside the corn seeds. Pioneer 37R71 corn was planted on May 11, 1998 at a rate of 62,000 seeds/ha. Each plot was 10.6 m long and consisted of 4 rows spaced 76 cm apart.

The tillage experiments were conducted as a randomized complete block design in a factorial arrangement with three replications. The factors consisted of herbicide application timing and herbicide treatment. Herbicides were applied when the corn was at the 2-leaf and 4-leaf stages. Corn leaf stages are described as the number of visible leaves. Herbicide treatments included isoxaflutole alone (BALANCE) at 105 g/ha and in combination with 1.1 kg/ha of metolachlor/benoxacor (DUAL II). Each of these treatments were applied either alone or with either SMS or AFS-MTMS. Each repellent was used at a rate of 0.5% v/v. Additional treatments not included in the factorial arrangement were an untreated check and a weed-free check. All herbicides were applied with a tractor mounted, compressed-air sprayer calibrated to deliver 2.6 L/ha at 207 kPa using 8003 E flat-fan

nozzles.

Corn tolerance was evaluated 30 days after planting (DAP) by visually evaluating plants for bleaching and necrotic symptoms and also by measuring corn height (base of plant to the crown) 40 DAP. Weed control by species was visually evaluated 60 DAP. Visual evaluations were based on a scale of 0 (no effect) to 100% (complete weed or crop death). Corn grain yield was determined by harvesting the center two rows of each plot with a plot combine. Seed weight was adjusted to 15% moisture.

Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at  $\alpha=0.05$ . Data were combined over years when treatment and/or application timing by year interactions were not significant  $\alpha=0.05$ . Non-transformed means for corn injury and weed control are presented since arcsine and square root transformations did not alter the interpretation of the data. Corn height and yield results were converted to a percent of the weed-free treatment after separation.

Herbicide application times, corn stages, weed heights, and densities for the field trial in 1998 are presented in Table 4 and rainfall data in Table 5.

Table 4

	1998	
	2-Leaf <sup>a</sup>	4-Leaf
Days after planting <sup>b</sup>	9	15
Corn		
Leaves with collars	1	2
Ave height (cm)	10	13
Giant foxtail		
Ave. height (cm)	0.6	4
Density (plants/m <sup>2</sup> )	33	65
Broadleaf weeds <sup>c</sup>		
Ave. height	0.6	3
Ave density (plants/m <sup>2</sup> )	44	44

<sup>a</sup> Corn leaf stage refers to the number of visible leaves.

<sup>b</sup> Corn planted May 11, 1998

<sup>c</sup> Broadleaf weeds include: common lambsquarters, redroot pigweed, common ragweed, and velvetleaf.



Table 5

Days after planting	Amount of Rainfall
	1998
	-----mm-----
0-7	2
8-14	3
15-21	12
22-28	0
Total	17

Corn (ZEAMX) in the 2-leaf or 4-leaf stage was treated with herbicidal compositions consisting of BALANCE (isoxaflutole); BALANCE and DUAL II (metolachlor and the safener benoxacor); BALANCE and SMS; BALANCE and AFS-MTMS; BALANCE, DUAL II and SMS; or BALANCE, DUAL, and AFS-MTMS. The percent injury was determined 6 DAT, 12 DAT, and 30 DAP. As weed controls, the annual grasses (ANGR) and common lambsquarters, *Chenopodium album* L. (CHEAL) were treated with herbicidal compositions consisting of BALANCE; BALANCE and DUAL II; BALANCE and SMS; BALANCE and AFS-MTMS; BALANCE, DUAL II and SMS; or BALANCE, DUAL, and AFS-MTMS. The percent injury for the weed controls was determined 30 DAP.

The results are shown in Table 6. The results show that the repellent adjuvants SMS and AFS-MTMS were effective in reducing the percent injury to corn caused by the herbicide when either was included in herbicidal compositions consisting of BALANCE and DUAL II. AFS-MTMS was particularly effective, when it was included in the herbicidal composition and applied to 2-leaf corn plants, corn injury 6 DAT was only 7.3%, whereas without AFS-MTMS, the corn injury was 65%. By 12 DAT and beyond, corn injury of caused by the BALANCE and DUAL II composition containing AFS-MTMS was not detectable whereas without AFS-MTMS, the injury remained about 55-50%. The weed controls show that SMS and AFS-MTMS do not appear to reduce the efficacy of BALANCE and DUAL II

to control weeds.

The results in Table 6 also show that SMS and AFS-MTMS reduced injury to corn at the 4-leaf stage when included in herbicidal compositions containing both BALANCE and DUAL II. In particular, the percent injury to corn was significantly reduced when the herbicidal composition included AFS-MTMS as the repellent adjuvant. The results further show that including a safener in the herbicide composition (the benoxacor) had no safening effect when used in combination with isoxaflutole. These results demonstrate that the present invention is useful and effective under actual field conditions.

5.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200

Table 6

Weed Code: Crop Code:						ZEAMX			ANGR	CHEAL
Treatment/Evaluation Interval Date evaluated						6 DAT 5-26-98	12 DAT 6-1-98	30 DAP 6-9-98	30 DAP 6-9-98	30 DAP 6-9-98
No.	Composition	Form Amt	Rate	Rate Unit	Grow Stg	Injury Percent			Control Percent	
1	BALANCE	75	1.5	OZ A/A	2-leaf	0.0	0.0	0.0	41.7	98.3
2	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	0.0	0.0	0.0	50.0	100.0
3	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	0.0	0.0	0.0	35.0	97.3
4	BALANCE DUAL II	75 7.8	1.5	OZ A/A LB A/A	2-leaf	65.0	55.0	50.0	99.3	100.0
5	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	23.3	5.0	5.7	96.0	100.0
6	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	7.3	0.0	0.0	98.7	100.0
7	BALANCE	75	1.5	OZ A/A	4-leaf		16.7	9.7	48.3	100.0
8	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	4-leaf		15.0	10.7	41.7	100.0
9	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	4-leaf		10.0	10.0	55.0	48.3
10	BALANCE DUAL II	75 7.8	1.5 1.0	OZ A/A LB A/A	4-leaf		81.7	83.3	100.0	100.0
11	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf		48.3	30.0	100.0	99.3
12	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf		26.7	16.7	98.7	99.7
13	Untreated						0.0	0.0	0.0	0.0
LSD (P= 0.05)						4.65	7.99	7.52	14.50	15.82
Standard Deviation						2.61	4.74	4.46	8.61	9.39
CV						19.11	23.87	26.86	12.94	10.68

BALANCE - isoxaflutole; DUAL II - metolachlor/benoxacor ; DAT - day after treatment, DAP - day after planting; ZEAMX - corn, *Zea mays* L.; ANGR - annual grasses, CHEAL - common lambsquarter, *Chenopodium album* L.

Additional weed controls consisted of the weeds giant foxtail, *Setaria faberi* Herrm. (SETFA);

5

contained either the SMS or AFS-MTMS repellent adjuvant and herbicidal compositions that did not contain either repellent adjuvant.

Table 7

Weed Code:						AMARE	AMBEL	ABUTH
Treatment/Evaluation Interval Date evaluated						30 DAP 6-9-98	30 DAP 6-9-98	30 DAP 6-9-98
No.	Composition	Form Amt	Rate Rate	Unit	Grow Stg	Control Percent		
1	BALANCE	75	1.5	OZ A/A	2-leaf	100.0	100.0	98.3
2	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	100.0	100.0	99.3
3	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	95.0	100.0	100.0
4	BALANCE DUAL II	75 7.8	1.5	OZ A/A LB A/A	2-leaf	100.0	100.0	100.0
5	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	100.0	100.0	97.7
6	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	100.0	100.0	99.3
7	BALANCE	75	1.5	OZ A/A	4-leaf	100.0	100.0	95.7
8	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	4-leaf	100.0	100.0	97.0
9	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	4-leaf	98.3	97.3	95.7
10	BALANCE DUAL II	75 7.8	1.5 1.0	OZ A/A LB A/A	4-leaf	100.0	100.0	98.3
11	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf	100.0	100.0	97.0
12	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf	100.0	100.0	98.3
13	Untreated					0.0	0.0	0.0
LSD (P= 0.05)						4.16	2.16	2.89
Standard Deviation						2.47	1.28	1.72
CV						2.69	1.39	1.9

BALANCE - isoxaflutole, DUAL II - metolachlor/benoxacor, DAT - day after treatment; DAP - day after planting, AMARE -redroot pigweed, *Amaranthus retroflexus* L. , AMBEL - common ragweed, *Ambrosia artemisiifolia* L. , ABUTH - velvetleaf, *Abutilon theophrasti* Medicus

The results of Tables 6 and 7 are condensed in Table 8, which shows that the present invention prevented injury to corn caused by herbicide component

of the composition without substantially affecting the ability of the herbicide component to control weed growth. Additional data was added to Table 8. The additional data shows that adding SMS or AFS-MTMS to the herbicide mixture prevented the herbicide component from affecting the growth of the corn. Also new in Table 8 is data showing that the herbicide mixture containing either SMS or AFS-MTMS did not substantially alter the herbicide component's ability to control the growth of giant foxtail (SETFA). Significantly, the grain yield of corn treated with the present invention was greater than the yields of corn treated with the herbicide mixture lacking either SMS or AFS-MTMS or the untreated controls.

Therefore, the results of the field trial show that the present invention renders the herbicides comprising the invention safe for use on corn while not reducing the ability of the herbicides comprising the present invention to control a wide variety of weeds.

Table 8

Herbicide	Application Stage	Corn		Weed Control <sup>c</sup>				Grain Yield
		Injury <sup>a</sup>	Height <sup>b</sup>	SETFA	CHEAL	AMARE	ABUTH	
		—%—	—cm—	—%—				kg/ha
Isoxaflutole <sup>d</sup>	2-leaf	0	61	73	95	98	99	10420
Isoxaflutole + SMS <sup>e</sup>	2-leaf	0	61	72	96	98	100	10146
Isoxaflutole + AFS-MTMS	2-leaf	0	61	56	85	92	100	10146
Isoxaflutole + Metolachlor <sup>f</sup>	2-leaf	50	41	95	100	100	100	9617
Isoxaflutole + metolachlor + SMS	2-leaf	6	56	91	100	97	98	10277
Isoxaflutole + metolachlor + AFS-MTMS	2-leaf	0	62	86	98	100	97	10527
Isoxaflutole	4-leaf	10	61	70	96	100	100	9995
Isoxaflutole + SMS	4-leaf	11	61	63	100	100	100	9881
Isoxaflutole + AFS-MTMS	4-leaf	10	58	68	83	95	96	10728
Isoxaflutole + Metolachlor <sup>a</sup>	4-leaf	83	32	95	100	100	98	9192
Isoxaflutole + metolachlor + SMS	4-leaf	30	50	99	100	100	100	9508
Isoxaflutole + metolachlor + AFS-MTMS	4-leaf	17	58	96	100	100	100	9993
Untreated		0	64	0	0	0	0	8528
LSD <sub>50</sub>		8	7	13	9	8	3	850

<sup>a</sup> Corn injury evaluated 30 DAP which was 21 days after 2-leaf application and 15 days after 4-leaf application

<sup>b</sup> Corn height was measured 40 DAP.

<sup>c</sup> Weed control was evaluated 60 DAP.

<sup>d</sup> The rate of isoxaflutole was 105 g/ha

<sup>e</sup> All adjuvants were applied at 0.5% v/v.

<sup>f</sup> The rate of metolachlor was 1121 g/ha

<sup>g</sup> The formulation of metolachlor contained the herbicide safener benoxacor

EXAMPLE 3

This example was performed as in Example 1, except that adjuvants SMS and AFS-MTMS were added to an isoxaflutole tank-mixture that contained an acetochlor herbicide and the safener MON-13900. As in Example 1, the herbicide applications were made to 2-leaf corn, and the herbicide treatments consisted of isoxaflutole at 105 g/ha and isoxaflutole tank-mixed with 1.8 kg/ha of acetochlor/MON-13900 applied either alone or in combination with either NIS at 0.25% (v/v), SMS at 0.5% (v/v), or AFS-MTMS at 0.5% (v/v).

The herbicide compositions were applied through an 8003 E flat fan nozzle delivering 234 L/ha at a pressure of 172 kPa. Corn tolerance was evaluated 8 DAT by visually evaluating the plants for bleaching and necrotic symptoms and also by measuring corn height (base of the plant to its crown). Visual corn injury ratings were based on a scale from 0 to 100, with 0 indicating no effect and 100 indicating plant death. Corn height was measured in cm and presented as a percent of the non-treated plants, with 0 indicating total reduction in plant height and 100 indicating height equal to the non-treated plants. All experiments were conducted twice as completely randomized designs with four replications. Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at  $\alpha=0.05$ . Statistical analysis indicated no experimental run interactions, so the data were combined and reported as the means of two experiments. Non transformed means are presented since arcsine and square root transformations did not alter the interpretation of the data.

Table 9 shows the corn injury and height reductions as influenced by SMS or AFS-MTMS combined with isoxaflutole alone and in combination with acetochlor/MON-13900 when applied postemergence in the greenhouse. The table shows that the percent injury to



corn treated with isoxaflutole in combination with acetochlor/MON-13900 was reduced to 20% when SMS was added to the combination. When AFS-MTMS was added to the combination of isoxaflutole and acetochlor/MON-13900 there was no visible injury to the corn. The table also shows that isoxaflutole in combination with acetochlor/MON-13900 reduced corn height by about 50% whereas adding either SMS or AFS-MTMS to the composition reduced the effect of the composition on corn height. In particular, in the presence of SMS, the height of the corn was reduced by only 20% and in the presence of AFS-MTMS, the height of the corn was virtually unaffected.

Table 9

Treatment <sup>a</sup>	Rate	Injury <sup>b</sup>		Height	
		Isoxaflutole <sup>c</sup>	Isoxaflutole <sup>c</sup> + acetochlor <sup>d</sup>	Isoxaflutole	Isoxaflutole <sup>c</sup> + acetochlor <sup>d</sup>
		-----%	-----%	-----% of control-----	-----%
Alone		0	64	99	47
+NIS	0.25% v/v	13	68	88	45
+SMS	0.5% v/v	0	20	100	79
+AFS-MTMS	0.5% v/v	0	0	100	98
LSD <sub>0.05</sub>		-----4-----	-----8-----		

<sup>a</sup> Treatments were applied to 2-leaf (12) corn.

<sup>b</sup> Visual injury ratings and corn heights were evaluated 8 DAT.

<sup>c</sup> Isoxaflutole was applied at 105 g/ha.

<sup>d</sup> The formulation of acetochlor contained the herbicide safener MON-13900 and was applied at 1.1 kg/ha

#### EXAMPLE 4

This example was conducted to determine the effect of SMS and AFS-MTMS on foliar retention of isoxaflutole either alone or in combination with metolachlor/benoxacor. A version of the technique reported by Boldt and Putnam, Weed Science 28: 474-477 (1980) was used. Herbicide treatments examined were isoxaflutole applied alone at 105 g/ha and isoxaflutole tank-mixed with 1.1 kg/ha of metolachlor/benoxacor.

Each of these herbicide treatments were applied alone or with either SMS or AFS-MTMS. The repellent adjuvants were each used at a rate of 0.5% v/v. The spray treatments, which included CHICAGO SKY BLUE (a product available from Sigma Chemical Co., St. Louis, Missouri) at 2.5 g/L, were applied to 2-leaf corn. Immediately after application, the whole plant was harvested and rinsed with distilled water containing the non-ionic surfactant X-77 (a product available from Valent U.S.A. Corp., Walnut Creek, California) at 0.25% v/v. The absorbance of the rinsate was determined spectrophotometrically at 625 nm. Dye retention ( $\mu\text{g}/\text{plant}$ ) was calculated from a standard curve.

Table 10 shows the spray retention of isoxaflutole as influenced by either SMS or AFS-MTMS, alone and in combination with metolachlor/benoxacor applied to 2-leaf corn in the greenhouse. The results show that neither SMS or AFS-MTMS increased the retention of isoxaflutole alone by the corn plant. The results further show that AFS-MTMS was particularly effective in reducing isoxaflutole retention when the composition further included metolachlor/benoxacor.

Table 10

Treatment	Rate	Spray retention	
		Isoxaflutole <sup>a</sup>	Isoxaflutole + metolachlor <sup>b</sup>
-----µg of isoxaflutole/plant-----			
Alone		4.4	15.1
+SMS	0.5% v/v	5.8	17.3
+AFS-MTMS	0.5% v/v	4.4	6.2
LSD <sub>0.05</sub>		-----2.2-----	

<sup>a</sup> rates; isoxaflutole at 105 g/ha, metolachlor/benoxacor at 1.1 kg/ha.

<sup>b</sup> The formulation contained the herbicide safener benoxacor

EXAMPLE 5

This example was to determine whether the adjuvants NIS, MTMS, SMS, AFS-MTMS, and PEAS affected weed control from early postemergence applications of isoxaflutole and isoxaflutole tank-mixed with metolachlor/benoxacor. Reduced rates of isoxaflutole (53 g/ha) and isoxaflutole tank-mixed with metolachlor/benoxacor (0.55 kg/ha) were applied alone and with either of NIS, MTMS, SMS, AFS-MTMS, or PEAS to 2-leaf velvetleaf (ABUTH) (3.5 to 5 cm) and 3-leaf barnyardgrass (5 to 10 cm). Velvetleaf and barnyardgrass controls were evaluated 21 DAT and shoots of both species were harvested to measure dry weight per pot. The results of this example are consistent with the weed control results presented in Tables 6, 7 and 8 of Example 2, which showed that the present invention was effective against weeds.

While the present invention is described herein with reference to illustrated embodiments, it should be understood that the invention is not limited hereto. Those having ordinary skill in the art and access to the teachings herein will recognize additional modifications and embodiments within the scope thereof. Therefore, the present invention is limited only by the Claims attached herein.